

CLAIMS

1. A reagent comprising:
a cytokine or chemokine-derived peptide or a cytokine-receptor-derived
5 peptide or chemokine-receptor derived-peptide;
a carrier protein; and
an adjuvant.
2. The reagent according to claim 1 wherein the cytokine or chemokine
is selected from the group consisting of IL-4, IL-13, IL-5, IL-9, IL-25, eotaxin or
10 TARC 3.
3. The reagent according to claim 1 wherein the cytokine is IL-4 and the
peptide is selected from the group consisting of: 6 or more consecutive residues of
DITLQEIIKTLNSLT (SEQ ID No. 1, amino acids 4-18 of IL-4); 6 or more
consecutive residues of EKETFCAATVLRQFYSHH (SEQ ID No. 2, amino acids
15 41-59 of IL-4); 6 or more consecutive residues of
QQFHRHKQLIRFLKRLDRNLWGLA (SEQ ID No. 3, amino acids 71-94 of IL-4); 6
or more consecutive residues of TLENFLERLKTIMREKYS (SEQ ID No. 4, amino
acids 108-125 of IL-4); 6 or more consecutive residues of EQKTLCTELTVTDIFA
(SEQ ID No. 5, amino acids 19-34 of IL-4); and 6 or more consecutive residues of
20 AGLNSCPVKE (SEQ ID No. 6, amino acids 94-103 of IL-4).
4. The reagent according to claim 1 wherein the cytokine is IL-13 and
the peptide is selected from the group consisting of 6 or more consecutive
residues of VPPSTALRELIEELVNITQ (SEQ ID No. 7, amino acids 4-22 of IL-13);
6 or more consecutive residues of MYCAALESII (SEQ ID No. 8, amino acids 43-
25 52 of IL-13); 6 or more consecutive residues of VAQFVKDLLLHLKK (SEQ ID No.
9, amino acids 92-105 of IL-13); 6 or more consecutive residues of
KDLLLHLKKLFREGRFN (SEQ ID No. 10, amino acids 97-113 of IL-13); 6 or more
consecutive residues of KIEVAQFVKDLLLHLKKLFREGRFN (SEQ ID No. 11,
amino acids 89-113 of IL-13); 6 or more consecutive residues of
30 SAIEKTQRMLSGFC (SEQ ID No. 12, amino acids 58-71 of IL-13); and 6 or more
consecutive residues of NVSGCSAIEKTQRMLSGFC (SEQ ID No. 13, amino acids
53-71 of IL-13).
5. The reagent according to claim 1 wherein the cytokine is IL-9 and the

BEST AVAILABLE COPY

peptide is selected from the group consisting of 6 or more consecutive residues of PTLAGILDINF (SEQ ID No. 14, amino acids 4-11 of IL-9); 6 or more consecutive residues of TRYPLIFSRVKKSVE (SEQ ID No. 15, amino acids 65-79 of IL-9); 6 or more consecutive residues of NALTFLKSLLEI (SEQ ID No. 16, amino acids 102-113 of IL-9); 6 or more consecutive residues of PASKCHCSANVTSCCLCLG (SEQ ID No. 17, amino acids 23-40 of IL-9); 6 or more consecutive residues of CTRPCFSE (SEQ ID No. 18, amino acids 46-53 of IL-9); and 6 or more consecutive residues of KNNKCPYFSCEQPCN (SEQ ID No. 19, amino acids 82-96 of IL-9).

6. The reagent according to claim 1 wherein the cytokine is IL-5 and the peptide is selected from the group consisting of 6 or more consecutive residues of PTSALVKETLALLSTHRTLLIA (SEQ ID No. 20, amino acids 6-27 of IL-5); 6 or more consecutive residues of PTSALVKETLALLST (SEQ ID No. 21, amino acids 6-20 of IL-5); 6 or more consecutive residues of HRTLLIA (SEQ ID No. 22, amino acids 21-27 of IL-5); 6 or more consecutive residues of EERRRVNQFLD (SEQ ID No. 23, amino acids 88-98 of IL-5); 6 or more consecutive residues of TVERLFKNLSLIKK (SEQ ID No. 24, amino acids 64-77 of IL-5); and PVHKNH (SEQ ID No. 25, amino acids 36-41 of IL-5).

7. The reagent according to claim 1 wherein the cytokine receptor is IL-4 α receptor and the peptide is selected from the group consisting of 6 or more consecutive residues of LYQLVFLLSEAH (SEQ ID No. 26, amino acids 36-47 of IL-4 α receptor); 6 or more consecutive residues of LLMDDVVSAD (SEQ ID No. 27, amino acids 63-72 of IL-4 α receptor); 6 or more consecutive residues of PPDNYLYNH (SEQ ID No. 28, amino acids 123-131 of IL-4 α receptor); and 6 or more consecutive residues of WAQAYNTT (SEQ ID No. 29, amino acids 177-186 of IL-4 α receptor).

8. The reagent according to claim 1 wherein the adjuvant is selected from the group consisting of: CpG oligodeoxynucleotides, alum, novasomes and liposomes.

9. The reagent according to claim 1 wherein the carrier protein is a highly immunogenic carrier protein.

10. The reagent according to claim 9 wherein the carrier protein is

BEST AVAILABLE COPY

selected from the group consisting of hepatitis B core antigen and hepatitis B surface antigen.

11. The reagent according to claim 10 wherein the peptide is fused into an immunodominant region of the carrier protein.

5 12. A method of inducing an immune response in an individual, comprising:
administering to an individual in need of such a treatment, an effective amount of a composition comprising:

10 a cytokine-derived peptide or chemokine-derived peptide or a cytokine-receptor derived peptide or chemokine-receptor derived-peptide;
a carrier protein; and
an adjuvant.

13. The method according to claim 12 wherein the cytokine is selected from the group consisting of IL-4, IL-13, IL-5, IL-9 or IL-25.

15 14. The method according to claim 12 wherein the cytokine is IL-4 and the peptide is selected from the group consisting of: 6 or more consecutive residues of DITLQEIIKTLNSLT (SEQ ID No. 1, amino acids 4-18 of IL-4); 6 or more consecutive residues of EKETFCRAATVLRQFYSHH (SEQ ID No. 2, amino acids 41-59 of IL-4); 6 or more consecutive residues of
20 QQFHRHKQLIRFLKRLDRNLWGLA (SEQ ID No. 3, amino acids 71-94 of IL-4); 6 or more consecutive residues of TLENFLERLKTIMREKYS (SEQ ID No. 4, amino acids 108-125 of IL-4); 6 or more consecutive residues of EQKTLCTELTVTDIFA (SEQ ID No. 5, amino acids 19-34 of IL-4); and 6 or more consecutive residues of
AGLNSCPVE (SEQ ID No. 6, amino acids 94-103 of IL-4).

25 15. The method according to claim 12 wherein the cytokine is IL-13 and the peptide is selected from the group consisting of 6 or more consecutive residues of VPPSTALRELIEELVNITQ (SEQ ID No. 7, amino acids 4-22 of IL-13); 6 or more consecutive residues of MYCAALES LI (SEQ ID No. 8, amino acids 43-52 of IL-13); 6 or more consecutive residues of VAQFVKDLLLLHLKK (SEQ ID No.
30 9, amino acids 92-105 of IL-13); 6 or more consecutive residues of KDLLLHLKKLFREGRFN (SEQ ID No. 10, amino acids 97-113 of IL-13); 6 or more consecutive residues of KIEVAQFVKDLLLLHLKKLFREGRFN (SEQ ID No. 11, amino acids 89-113 of IL-13); 6 or more consecutive residues of

BEST AVAILABLE COPY

SAIEKTQRMLSGFC (SEQ ID No. 12, amino acids 58-71 of IL-13); and 6 or more consecutive residues of NVSGCSAIEKTQRMLSGFC (SEQ ID No. 13, amino acids 53-71 of IL-13).

16. The method according to claim 12 wherein the cytokine is IL-9 and
5 the peptide is selected from the group consisting of 6 or more consecutive residues of PTLAGILDINF (SEQ ID No. 14, amino acids 4-11 of IL-9); 6 or more consecutive residues of TRYPLIFSRVKKSVE (SEQ ID No. 15, amino acids 65-79 of IL-9); 6 or more consecutive residues of NALTFLKSLLEI (SEQ ID No. 16, amino acids 102-113 of IL-9); 6 or more consecutive residues of
10 PASKCHCSANVTSCCLCG (SEQ ID No. 17, amino acids 23-40 of IL-9); 6 or more consecutive residues of CTRPCFSE (SEQ ID No. 18, amino acids 46-53 of IL-9); and 6 or more consecutive residues of KNNKCPYFSCEQPCN (SEQ ID No. 19, amino acids 82-96 of IL-9).

17. The method according to claim 12 wherein the cytokine is IL-5 and
15 the peptide is selected from the group consisting of 6 or more consecutive residues of PTSALVKETLALLSTHRTLLIA (SEQ ID No. 20, amino acids 6-27 of IL-5); 6 or more consecutive residues of PTSALVKETLALLST (SEQ ID No. 21, amino acids 6-20 of IL-5); 6 or more consecutive residues of HRTLLIA (SEQ ID No. 22, amino acids 21-27 of IL-5); 6 or more consecutive residues of EERRRVNQFLD
20 (SEQ ID No. 23, amino acids 88-98 of IL-5); 6 or more consecutive residues of TVERLFKNLSLIKK (SEQ ID No. 24, amino acids 64-77 of IL-5); and PVHKNH (SEQ ID No. 25, amino acids 36-41 of IL-5).

18. The method according to claim 12 wherein the cytokine receptor is IL-4 α receptor and the peptide is selected from the group consisting of 6 or more
25 consecutive residues of LYQLVFLLSEAH (SEQ ID No. 26, amino acids 36-47 of IL-4 α receptor); 6 or more consecutive residues of LLMDDVVSAD (SEQ ID No. 27, amino acids 63-72 of IL-4 α receptor); 6 or more consecutive residues of PPDNYLYNH (SEQ ID No. 28, amino acids 123-131 of IL-4 α receptor); and 6 or more consecutive residues of WAQAYNTT (SEQ ID No. 29, amino acids 177-186
30 of IL-4 α receptor).

19. The method according to claim 12 wherein the composition includes an adjuvant.

BEST AVAILABLE COPY

20. The method according to claim 19 wherein the adjuvant is selected from the group consisting of: CpG oligodeoxynucleotides, alum, novasomes and liposomes.

21. The method according to claim 12 wherein the carrier protein is a highly immunogenic carrier protein.

22. The method according to claim 21 wherein the carrier protein is selected from the group consisting of hepatitis B core antigen and hepatitis B surface antigen.

23. The method according to claim 22 wherein the peptide is fused into an immunodominant region of the carrier protein.

24. A method of treating, ameliorating or preventing asthma comprising: administering to an individual in need of such a treatment, an effective amount of a composition comprising:

a cytokine-derived peptide or a chemokine-derived peptide or a cytokine-receptor derived peptide or chemokine-receptor derived-peptide;

a carrier protein; and

an adjuvant.

25. The method according to claim 24 wherein the cytokine is selected from the group consisting of IL-4, IL-13, IL-2, IL-3, IL-5, IL-9 or IL-25.

26. The method according to claim 24 wherein the cytokine is IL-4 and the peptide is selected from the group consisting of: 6 or more consecutive residues of DITLQEIIKTLNSLT (SEQ ID No. 1, amino acids 4-18 of IL-4); 6 or more consecutive residues of EKETFCRAATVLRQFYSHH (SEQ ID No. 2, amino acids 41-59 of IL-4); 6 or more consecutive residues of QQFHRHKQLIRFLKRLDRNLWGLA (SEQ ID No. 3, amino acids 71-94 of IL-4); 6 or more consecutive residues of TLENFLERLKTIMREKYS (SEQ ID No. 4, amino acids 108-125 of IL-4); 6 or more consecutive residues of EQKTLCTELTVTDIFA (SEQ ID No. 5, amino acids 19-34 of IL-4); and 6 or more consecutive residues of AGLNSCPVKE (SEQ ID No. 6, amino acids 94-103 of IL-4).

27. The method according to claim 24 wherein the cytokine is IL-13 and the peptide is selected from the group consisting of 6 or more consecutive residues of VPPSTALRELIEELVNITQ (SEQ ID No. 7, amino acids 4-22 of IL-13); 6 or more consecutive residues of MYCAALES LI (SEQ ID No. 8, amino acids 43-

BEST AVAILABLE COPY

52 of IL-13); 6 or more consecutive residues of VAQFVKDLLLHLKK (SEQ ID No. 9, amino acids 92-105 of IL-13); 6 or more consecutive residues of KDLLLHLKKLFREGRFN (SEQ ID No. 10, amino acids 97-113 of IL-13); 6 or more consecutive residues of KIEVAQFVKDLLLHLKKLFREGRFN (SEQ ID No. 11, amino acids 89-113 of IL-13); 6 or more consecutive residues of SAIKTQRMLSGFC (SEQ ID No. 12, amino acids 58-71 of IL-13); and 6 or more consecutive residues of NVSGCSAIKTQRMLSGFC (SEQ ID No. 13, amino acids 53-71 of IL-13).

28. The method according to claim 24 wherein the cytokine is IL-9 and the peptide is selected from the group consisting of 6 or more consecutive residues of PTLAGILDINF (SEQ ID No. 14, amino acids 4-11 of IL-9); 6 or more consecutive residues of TRYPLIFSRVKKSVE (SEQ ID No. 15, amino acids 65-79 of IL-9); 6 or more consecutive residues of NALTFLKSLEI (SEQ ID No. 16, amino acids 102-113 of IL-9); 6 or more consecutive residues of PASKCHCSANVTSCCLG (SEQ ID No. 17, amino acids 23-40 of IL-9); 6 or more consecutive residues of CTRPCFSE (SEQ ID No. 18, amino acids 46-53 of IL-9); and 6 or more consecutive residues of KNNKCPYFSCEQPCN (SEQ ID No. 19, amino acids 82-96 of IL-9).

29. The method according to claim 24 wherein the cytokine is IL-5 and the peptide is selected from the group consisting of 6 or more consecutive residues of PTSALVKETLALLSTHRTLLIA (SEQ ID No. 20, amino acids 6-27 of IL-5); 6 or more consecutive residues of PTSALVKETLALLST (SEQ ID No. 21, amino acids 6-20 of IL-5); 6 or more consecutive residues of HRTLLIA (SEQ ID No. 22, amino acids 21-27 of IL-5); 6 or more consecutive residues of EERRRVNQFLD (SEQ ID No. 23, amino acids 88-98 of IL-5); 6 or more consecutive residues of TVERLFKNLSLIKK (SEQ ID No. 24, amino acids 64-77 of IL-5); and PVHKNH (SEQ ID No. 25, amino acids 36-41 of IL-5).

30. The method according to claim 24 wherein the cytokine receptor is IL-4 α receptor and the peptide is selected from the group consisting of 6 or more consecutive residues of LYQLVFLLEAH (SEQ ID No. 26, amino acids 36-47 of IL-4 α receptor); 6 or more consecutive residues of LLMDDVVSAD (SEQ ID No. 27, amino acids 63-72 of IL-4 α receptor); 6 or more consecutive residues of

BEST AVAILABLE COPY

PPDNYLYNH (SEQ ID No. 28, amino acids 123-131 of IL-4 α receptor); and 6 or more consecutive residues of WAQAYNTT (SEQ ID No. 29, amino acids 177-186 of IL-4 α receptor).

31. The method according to claim 24 wherein the composition includes
5 an adjuvant.

32. The method according to claim 31 wherein the adjuvant is selected from the group consisting of: CpG oligodeoxynucleotides, alum, novasomes and liposomes.

33. The method according to claim 24 wherein the carrier protein is a
10 highly immunogenic carrier protein.

34. The method according to claim 33 wherein the carrier protein is selected from the group consisting of hepatitis B core antigen and hepatitis B surface antigen.

35. The method according to claim 24 wherein the peptide is fused into
15 an immunodominant region of the carrier protein.

36. An expression system comprising:

a nucleic acid molecule deduced from a peptide selected from the group consisting of: 6 or more consecutive residues of DITLQEIIKTLNSLT (SEQ ID No. 1, amino acids 4-18 of IL-4); 6 or more consecutive residues of
20 EKETFCRAATVLRQFYSHH (SEQ ID No. 2, amino acids 41-59 of IL-4); 6 or more consecutive residues of QQFHRHKQLIRFLKRLDRNLWGLA (SEQ ID No. 3, amino acids 71-94); 6 or more consecutive residues of TLENFLERLKTIMREKYS (SEQ ID No. 4, amino acids 108-125); 6 or more consecutive residues of
25 EQKTLCTELTVTDIFA (SEQ ID No. 5, amino acids 19-34); 6 or more consecutive residues of AGLNSCPVKE (SEQ ID No. 6, amino acids 94-103 of IL-4); 6 or more consecutive residues of VPPSTALRELIEELVNITQ (SEQ ID No. 7, amino acids 4-22 of IL-13); 6 or more consecutive residues of MYCAALESII (SEQ ID No. 8, amino acids 43-52 of IL-13); 6 or more consecutive residues of
30 VAQFVKDLLLHLKK (SEQ ID No. 9, amino acids 92-105 of IL-13); 6 or more consecutive residues of KDLLLHLKKLFREGRFN (SEQ ID No. 10, amino acids 97-113 of IL-13); 6 or more consecutive residues of KIEVAQFVKDLLLHLKKLFREGRFN (SEQ ID No. 11, amino acids 89-113 of IL-13); 6 or more consecutive residues of SAIEKTQRMLSGFC (SEQ ID No. 12,

amino acids 58-71 of IL-13); 6 or more consecutive residues of NVSGCSAIEKTQRMLSGFC (SEQ ID No. 13, amino acids 53-71 of IL-13); 6 or more consecutive residues of PTLAGILDINF (SEQ ID No. 14, amino acids 4-11 of IL-9); 6 or more consecutive residues of TRYPLIFSRVKKSVE (SEQ ID No. 15, amino acids 65-79 of IL-9); 6 or more consecutive residues of NALTFLKSLLEI (SEQ ID No. 16, amino acids 102-113 of IL-9); 6 or more consecutive residues of PASKCHCSANVTSCCLCLG (SEQ ID No. 17, amino acids 23-40 of IL-9); 6 or more consecutive residues of CTRPCFSE (SEQ ID No. 18, amino acids 46-53 of IL-9); 6 or more consecutive residues of KNNKCPYFSCEQPCN (SEQ ID No. 19, amino acids 82-96 of IL-9); 6 or more consecutive residues of PTSALVKETLALLSTHRTLLIA (SEQ ID No. 20, amino acids 6-27 of IL-5); 6 or more consecutive residues of PTSALVKETLALLST (SEQ ID No. 21, amino acids 6-20 of IL-5); 6 or more consecutive residues of HRTLLIA (SEQ ID No. 22, amino acids 21-27 of IL-5); 6 or more consecutive residues of EERRRVNQFLD (SEQ ID No. 23, amino acids 88-98 of IL-5); 6 or more consecutive residues of TVERLFKNLSLIKK (SEQ ID No. 24, amino acids 64-77 of IL-5); PVHKNH (SEQ ID No. 25, amino acids 36-41 of IL-5); 6 or more consecutive residues of LYQLVFLLEAH (SEQ ID No. 26, amino acids 36-47 of IL-4 α receptor); 6 or more consecutive residues of LLMDVVVSAD (SEQ ID No. 27, amino acids 63-72 of IL-4 α receptor); 6 or more consecutive residues of PPDNYLYNH (SEQ ID No. 28, amino acids 123-131 of IL-4 α receptor); 6 or more consecutive residues of WAQAYNTT (SEQ ID No. 29, amino acids 177-186 of IL-4 α receptor); 6 or more consecutive residues of GPASVPTTCCFNLA (SEQ ID No. 30, amino acids 1-14 of eotaxin); 6 or more consecutive residues of FNLANRKIPLQRLES (SEQ ID No. 31, amino acids 11-25 of eotaxin); 6 or more consecutive residues of RITSGKCPQKAVIFKT (SEQ ID No. 32, amino acids 30-43 of eotaxin); and 6 or more consecutive residues of IFKTKLAKDICAD (SEQ ID No. 33, amino acids 40-52),

genetically fused to a nucleic acid molecule encoding a carrier protein.

30 37. The expression system according to claim 36 wherein the carrier protein is a highly immunogenic carrier protein.

38. The expression system according to claim 37 wherein the carrier

BEST AVAILABLE COPY

protein is selected from the group consisting of hepatitis B core antigen and hepatitis B surface antigen.

39. The expression system according to claim 38 wherein the peptide is fused into an immunodominant region of the carrier protein.

5 40. The reagent according to claim 1 wherein the chemokine is eotaxin and the peptide is selected from the group consisting of 6 or more consecutive residues of GPASVPTTCCFNLA (SEQ ID No. 30, amino acids 1-14 of eotaxin); 6 or more consecutive residues of FNLANRKIPLQRLES (SEQ ID No. 31, amino acids 11-25 of eotaxin); 6 or more consecutive residues of RITSGKCPQKAVIFKT
10 (SEQ ID No. 32, amino acids 30-43 of eotaxin); and 6 or more consecutive residues of IFKTKLAKDICAD (SEQ ID No. 33, amino acids 40-52).

41. The method according to claim 12 wherein the chemokine is eotaxin and the peptide is selected from the group consisting of 6 or more consecutive residues of GPASVPTTCCFNLA (SEQ ID No. 30, amino acids 1-14 of eotaxin); 6
15 or more consecutive residues of FNLANRKIPLQRLES (SEQ ID No. 31, amino acids 11-25 of eotaxin); 6 or more consecutive residues of RITSGKCPQKAVIFKT (SEQ ID No. 32, amino acids 30-43 of eotaxin); and 6 or more consecutive residues of IFKTKLAKDICAD (SEQ ID No. 33, amino acids 40-52).

42. The method according to claim 24 wherein the chemokine is eotaxin
20 and the peptide is selected from the group consisting of 6 or more consecutive residues of GPASVPTTCCFNLA (SEQ ID No. 30, amino acids 1-14 of eotaxin); 6 or more consecutive residues of FNLANRKIPLQRLES (SEQ ID No. 31, amino acids 11-25 of eotaxin); 6 or more consecutive residues of RITSGKCPQKAVIFKT (SEQ ID No. 32, amino acids 30-43 of eotaxin); and 6 or more consecutive
25 residues of IFKTKLAKDICAD (SEQ ID No. 33, amino acids 40-52).

BEST AVAILABLE COPY